

## Case Report

### Pseudoaneurysm of interventricular membranous septum: a rare late complication after repeated valve replacements

Ferhat C. Piskin<sup>1</sup>, Rabia Akilli<sup>2</sup>, Umur A. Pehlivan<sup>1</sup>, Erol H. Aksungur<sup>1</sup>, Kaigeldy Aikimbaev<sup>1</sup>

<sup>1</sup>Cukurova University Medical Faculty Radiology Department, Adana, Turkey

<sup>2</sup>Cukurova University Medical Faculty Cardiology Department, Adana, Turkey

#### Abstract

A 39-year-old female patient who had aortic and mitral valve replacement due to rheumatic heart disease was also found to have a cystic mass in the right ventricle on echocardiography during late routine control. The cystic mass identified on echocardiography was shown to be a pseudoaneurysm that develops from the membranous section of the interventricular septum and grows toward the right heart by ECG-triggered cardiac computed tomography performed for differential diagnosis. Because the patient did not accept the surgery, regular follow-ups were recommended.

**Key words:** pseudoaneurysm, membranous part of interventricular septum, cardiac computed tomography, aortic valve replacement, mitral valve replacement, cardiac surgery procedures, complication

(Heart, Vessels and Transplantation 2019; 3: doi: 10.24969/hvt.2019.147)

#### Introduction

Aneurysms of the membranous section of the interventricular septum (IVS) are rare pathologies (1). Aneurysms of this part of the septum are congenital and are in the form of true aneurysms, but a small number of aneurysms have been identified in the literature as pseudoaneurysms (2, 3).

The aim of this case report is to present a very rare case of developing of the pseudoaneurysm in the membranous part of IVS as a late complication after aortic and mitral valve replacement.

#### Case report

The asymptomatic 39-year-old female patient who had a cystic mass in the right heart identified by

echocardiography was evaluated by ECG-triggered cardiac computed tomography (CT) in our clinic.

In the clinical history of the patient, stenosis of the mitral valve due to rheumatic heart disease and the development of aortic valve insufficiency were treated with metallic prosthetic valve implantation (once on the aortic valve and twice on the mitral valve) in 2004.

In 2018, echocardiography identified a pulsatile cystic mass in the right atrium and right ventricle. This newly identified lesion had not been detected in the patient's previous echocardiography control. Physical examination and laboratory findings were normal, sinus rhythm was present on the electrocardiogram.

**Address for Correspondence:** Ferhat Can Piskin, Department of Radiology, Balcali Hospital, Medical Faculty, Cukurova University, Adana, Turkey; E-mail: ferhatcpiskin@gmail.com, Phone: +90 0555 736 8283, Fax: +90 322 338 6572

**Received:** 04.10.2019 **Revised:** 23.10.2019 **Accepted:** 28.20.2019

**Copyright © 2019 Heart, Vessels and Transplantation**

The patient was evaluated with ECG-triggered cardiac CT. The contrast agent was carried out to be two-phasic for the examination. A bolus injection of 80 mL of contrast with an iodine concentration of 370 mg I/mL was administered at 5.5 mL/s, followed by mixture (70% saline + 30% contrast agent) bolus of 5 mL/s by using a power injector with 18G intravenous needle in the right antecubital vein. The images were obtained with 160 detector CT devices (Aquilion; Toshiba Medical Systems, Tokyo, Japan) with a retrospective ECG trigger to include all cycles of the heart (systole and diastole). Then volume rendering (VR), maximum intensity projection (MIP), multiplanar reconstructions (MPR) and sine images were analyzed at the workstation (Vitrea; Vital Images Inc, Minnesota, USA). A pseudoaneurysm with a narrow neck, lobule and two separate segments in the membranous region of the IVS was detected at the short-axis (two chambers) and long-axis (four chambers) images of the left ventricle. The pseudoaneurysm's total size was 27x55 mm, and it was showing growth towards the right atrium, also replaced aorta and mitral metallic prosthetic valve were present (Fig. 1 A-C). The pseudoaneurysm lumen was empty and no thrombus formation was detected in it.

Surgical treatment was recommended but the patient did not accept this treatment due to the risks of complications. It was therefore decided that the patient should be followed up with anticoagulant treatment.

### Discussion

Aneurysms of the IVS membranous part are rare pathologies that account for 0.3% of congenital heart diseases. According to their etiology, they are divided into two types: congenital and acquired. Acquired IVS aneurysms are pseudoaneurysms, which develop secondary to surgical intervention (valve replacement) or trauma. Only two case reports have been published in the literature (2, 3).

IVS consists of two parts (membranous and muscular) with different histological characteristics. The membranous part of IVS consists of fibrotic tissue and is less resistant to pressure (5). In patients with mitral valve surgical replacement, the left ventricular wall can be damaged and separated from the fibrous skeleton during surgery. This can cause pseudoaneurysm in the subannular part of the left ventricle free wall (6). In our case, it was determined that a pseudoaneurysm developed in the IVS membranous septum, unlike the one expected. Although there is not enough literature on this subject, we estimate that this part of IVS was damaged during surgery in our patient with a history of aortic and mitral valve replacement and that the part of IVS was damaged due to the formation of fibrous tissue and pseudoaneurysm development in the long period.

Clinical signs of IVS aneurysms develop secondary to complications (pulmonary valve stenosis, thromboembolism and arrhythmia) that develop as a result of pressure or rupture of aneurysms (7). Echocardiography, contrast CT and magnetic resonance imaging (MRI) methods can be used as imaging methods for IVS aneurysms. Echocardiography has disadvantage due to the inability to provide the appropriate acoustic window for displaying the posterior structures of the heart. ECG-triggered cardiac CT and MRI allows better evaluation of anatomical details in 3-D high-resolution images. In addition, as recently introduced, dynamics and functions can be evaluated in phase contrast at ECG-triggered cardiac MRI (8).

Treatment methods of true aneurysms are chosen according to presence or absence of complications. In a patient in whom complications develop, surgical excision is applied and in other patient, anticoagulant treatment can be administered. Pseudoaneurysms have a higher risk of rupture because they do not contain a real wall structure, so the surgical treatment should be the first option for patient without considering the development of complications (8).




Figure 1. Contrast ECG-triggered cardiac CT images: A) Volume rendering, B) Short-axis MPR, C) Four-chamber MPR. The cystic lesion is observed in the membranous part of IVS. The cystic lesion develops into the right atrium and ventricle from left ventricle. In addition, the cystic lesion is projecting towards basis of the heart.  
CT- computed tomography, IVS- interventricular septum, MPR- multiplanar reconstruction

## Conclusion

Because the membranous part of IVS is a sensitive area for surgical interventions or traumas, pseudoaneurysm may develop in this part of IVS in patients with history of surgery at the long term. In addition, ECG-triggered cardiac CT or MRI should be used for the imaging of these anatomical structures.

**Peer-review:** External and internal

**Conflict of interest:** None to declare

**Authorship:** F.C.P., R.A., U.A.P., E.H.A., K.A. equally contributed to management of patient and preparation of case report

**Acknowledgement and funding:** None to declare

## References

1. Herpin D, Ciber MA, Boutaud P, Amiel A, Gaudeau B, Guillem JP, et al. Aneurysms of the interauricular septum. Review of the literature apropos of 8 cases. *Ann Cardiol Angeiol* 1986; 35: 35-42.
2. Agudo-Quilez P, Pozo E, Benedicto A, Reyes G, Olivera MJ, Caballero P, et al. Atrioventricular septum pseudoaneurysm as late complication after repeated mitral valve replacement. *Ann Thorac Surg.* 2017; 103: e55-6.
3. Yavuz S, Eris C, Goncu T, Sezen M, Ata Y, Turk T. An incidental aneurysm of the interventricular membranous septum. *Arch Iran Med* 2010; 13: 363-4.
4. Lazarev SM, Matsan VI. Aneurysm of the membranous heart septum. *Kardiologija* 1989; 29: 36-40.
5. Tardy MM, Galvaing G, Sakka L, Garcier JM, Chazal J, Filaire M. Embryology of the heart walls. *Morphol Bull Assoc Anat* 2013; 97: 2-11.
6. Sahan E, Sahan S, Karamanlioglu M, Gul M, Tufekcioglu O. Left ventricular pseudoaneurysm after mitral valve replacement: Review of pseudoaneurysms late after mitral valve replacement. *Herz* 2015; 40: 778-82.
7. Di Cesare E, Di Sibio A, Lanni G, Gennarelli A, Masciocchi C. Magnetic resonance imaging of AMS (Aneurysm of the Membranous Septum), review of the literature and case report. *J Radiol Case Rep* 2014; 8: 9-15.
8. Carcano C, Kanne JP, Kirsch J. Interventricular membranous septal aneurysm: CT and MR manifestations. *Insights Imaging* 2015; 7: 111-7.