

Relationship of preoperative NT-pro BNP with clinical, perioperative and prognostic markers in cardiac surgery: Preliminary study results

Irina A. Akhmedova, Taalaibek Z. Kudaiberdiev, Damirbek A. Abibillaev, Akylbek A. Zhooshev, Dolonbek E. Zaripov, Kaiyrnisa T. Tilemanbetova, Guliza N. Naizabekova

Scientific-Research Institute of Heart Surgery and Organ Transplantation, Bishkek, Kyrgyzstan

Abstract

Objective: We aimed to define the relationship of preoperative NT pro-BNP testing in routine practice of cardiac surgery with clinical, perioperative variables, surgical outcomes and complications in pediatric and adult cardiac surgery patients. In addition, we assessed relationship of NT pro-BNP with EuroSCORE II in adult patients undergoing cardiac surgery.

Methods: A total of 48 patients who underwent cardiac surgery in our institution were enrolled into study. According to age aspects pediatric (PG, n=20) and adult (AG, n=28) group of patients were separately evaluated. Each group further was categorized into two subgroups (group 1 and 2) on the basis of cut-off points of NT pro-BNP level (430 and 250pg/ml in adults and children, respectively). Preoperative NT-pro BNP was obtained from patients. Statistical tests were conducted to reveal differences in clinical and perioperative variables among NT-pro-BNP groups and relationship of the assay with baseline clinical and operative parameters, as well as postoperative outcomes.

Results: In adults undergoing cardiac surgery, statistically significant differences were found between groups with low and high levels of NT pro-BNP by NYHA classes (I to III, $p < 0.001$), body mass index (BMI) values (31.7(3.15) vs 25.9 (3.8) kg/cm², $p = 0.017$), estimated glomerular filtration rate (GFR_e) (111.4(26.4) vs 77.9(22.4) ml/min/1.73m² $p = 0.036$), moderately impaired renal function (20% vs 72.7%, $p = 0.049$), need for inotropes (28.6% vs 86.7%, $p = 0.006$), and mean EuroSCORE II (0.81 (0.19) vs 1.7 (0.5), $p = 0.008$). There was the positive significant correlation of NT pro-BNP levels with EuroSCORE II, NYHA class, need for inotropes, renal function and left ventricular (LV) function categories, while its negative correlation was found with BMI, LV ejection fraction and GFR_e parameters. In PG – NT pro-BNP was negatively correlated with age, body surface area, LV end-diastolic and end-systolic dimensions.

Conclusion: In adult patients undergoing cardiac surgery, preoperative NT-pro-BNP level correlated well with several baseline and clinical-operative parameters, including BMI, renal and LV function, NYHA class, need for inotropes support, and EuroSCORE II. Further studies are needed to define its exact diagnostic and prognostic significance in pediatric patients.

Key words: cardiac surgery procedures, congestive heart failure, NT-pro-BNP, correlation, prognosis, surgical risk, surgery outcomes, early postoperative complications, Euroscore, renal failure

(Heart Vessels Transplant 2020; 4; doi: 10.24969/hvt.2020.205)

Introduction

Heart failure may be a result of vast majority of cardiovascular disorders including myocardial damage, valve pathology, conduction and rhythm abnormalities, and pericardial diseases (1, 2). History of most of the patients reveals both cardiac and non-cardiac comorbid conditions as the underlying factors

of chronic heart failure (HF) (3). The incidence of HF is steadily increasing and remains the leading cause of hospitalization and in-hospital mortality in elderly population (age ≥ 65 years) (4-6). By the improvement of current management, great successes have been achieved both in pharmacologic and surgical treatment of chronic HF.

Address for Correspondence: Damirbek A. Abibillaev, Scientific Research Institute of Heart Surgery and Organ Transplantation, Bishkek, Kyrgyzstan E-mail: kg.damir.da@gmail.com

Received: 14.05.2020 **Revised:** 02.06.2020 **Accepted:** 05.06.2020

Copyright© 2020 Heart, Vessels and Transplantation

Heart surgeries are quite challenging procedures which require a cardiac-pulmonary bypass (CPB) and other settings. Additionally, high risk of complications is observed in postoperative period. Advanced age of patients and huge number of comorbid issues lead to increase of disease severity requiring more complex operations including simultaneous, hybrid and multistage procedures (7, 8). Furthermore, despite the significant improvement of healthcare, surgical operations performed in children with congenital heart diseases (CHD) are accompanied by high morbidity and mortality both in early and late post-op. Implementation of the robust prognostic biomarkers besides the routine ones used in daily clinical practice is crucial in postoperative management of pediatric patients in intensive care unit (9).

Several multifactorial scores have been used for pre-operative (pre-op) risk stratification and surgery outcomes. But, no single risk assessment tool for cardiac surgery includes pre-op measurement of brain-type natriuretic peptides (BNP) in spite of their accurate correlation with ventricular dysfunction and HF. (10) On the other hand, some studies recommend cautious application of risk assessment models in cardiac surgery for different population groups, due to existence of great variety by gender, ethnicity and geographic factors (11).

BNP, a valuable marker of HF is the member of natriuretic peptides family and synthesized, then secreted into bloodstream in response to volume overload or conditions causing myocardial stretching of ventricles. Compared to atrial natriuretic peptides (ANP), BNP is synthesized not only by atria, but also by ventricles, particularly in patients with HF (12). Circulating concentrations of several cardiac natriuretic peptides, including ANP, BNP and their N-terminal prohormones (NT-pro-ANP and NT-pro-BNP) are elevated both in patients with HF signs and asymptomatic left ventricular (LV) dysfunction (13-15). In practical sense, BNP and NT-pro-BNP are interchangeable and can be used according to local preference and availability. For diagnostic workup of mild HF and subclinical LV dysfunction NT-pro-BNP is preferred over BNP (15).

Duceppe E et al. proposed the prognostic role of pro-BNP in adverse cardiovascular outcomes and mortality risk assessment of patients undergoing non-cardiac surgery within post-operative (post-op) thirty days (16). In cardiac surgery, pre-op levels of NT pro-BNP well correlate with ventilation time, duration of intensive care stays and perioperative pharmacological inotropic support (17). According to Hutfless et al, the level of BNP>385pg/ml may predict

post-op complications and mortality within twelve months after the cardiac surgery (10).

The study of Liu H. et al. revealed the level of NT pro-BNP at 2773.5 pg/ml as a cut-off value (sensitivity 63.6% and specificity 80.8%) for mortality prognosis within 30 days after surgery. A mean NT pro-BNP level of cases with adverse clinical outcomes significantly exceeded the mean NT pro-BNP assays of uncomplicated patients (1998.5 pg/ml vs 630.5pg/ml, $p = 0.016$) (18). However, the evidence of clinical and prognostic utility of NTproBNP in cardiac surgery is still emerging.

In this preliminary study, we objected to define the relationship of preoperative NT pro-BNP testing in routine practice of cardiac surgery with clinical, perioperative variables (cardiopulmonary bypass time, aortic cross-clamp time, ventilation duration, need for inotropes, duration of intensive care stays), surgical outcomes and complications in pediatric and adult cardiac surgery patients. In addition, we assessed relationship of NT pro-BNP with EuroSCORE II in adult patients undergoing cardiac surgery.

Methods

Study design

Retrospective, observational cohort study.

Study population

A total of 48 patients enrolled into study underwent elective cardiac surgery in Scientific-Research Institute of Heart Surgery and Organ Transplantation (SRIHSOT) within two months of 2019 (September-November). Pediatric patients constituted 42% ($n=20$) operated due to CHD, adults comprised 58% ($n=28$) and underwent coronary artery bypass surgery (CABG), aortic (AVR) and mitral valve (MVR) replacement surgeries and myxoma resection surgery.

Study population was categorized into two groups by age aspects: pediatric (PG) and adult (AG) ones. Further each age group was divided into two subgroups, according to NT pro-BNP levels: 1st subgroup included cases with less than 430pg/ml (AG) (19) and 250pg/ml (PG) (20), 2nd subgroup more than those cut-off points.

The NYHA IV class patients were excluded, though they did not undergo surgery. Patients with altered renal function (chronic kidney disease IIIB-V) were also excluded. Due to absence of NT pro-BNP data 4 pediatric and 6 adult patients were excluded from the study.

Informed consent for procedures were obtained from all adult patients and from parents (guardians) of children patients.

Data analysis

We collected the following baseline demographic data from patient records: age (expressed in years (AG) and months (PG)), sex (male or female), body mass index (by Kettle formula in AG), body surface area (by Haycock formula in PG), underlying diagnoses, type of surgical operations, presence of risk factors for adverse outcomes of CHF (presence or absence of the dyslipidemia/atherosclerosis, myocardial infarction (MI), arterial hypertension, diabetes mellitus (DM), pre-op cardiac arrhythmia).

For pre-op assessment of CHF in adults we used New-York Heart Association functional classification (NYHA) proposed in 1964, while for pediatric patients Modified Ross classification for pediatric heart failure (21, 22).

For baseline clinical data we collected serum creatinine levels, estimated glomerular filtration rate (eGFR) assessed by modified Cocroft-Gault formula (23), renal function categories (normal renal function (≥ 85 ml/min/1.73m²), moderately or severely impaired (85-45 ml/min/1.73m² and < 45 ml/min/1.73m²), echocardiographic LV indices (LV end-diastolic (LVEDD) and end-systolic dimensions, (LVESD), LV ejection fraction (EF) by Teichholz and Simpson methods) and LV systolic function categories (reduced EF $<40\%$, moderate (EF 40-50%) or preserved (EF $\geq 50\%$)), Doppler-derived systolic pulmonary artery pressure (SPAP), pulmonary artery hypertension (PAH) categories (normal (<30 mmHg), moderate PAH (31-55 mmHg) and severe (>55 mmHg)).

For perioperative data we collected cardiac-pulmonary bypass (CPB) and aortic cross-clamp (ACCT) times, duration of ICU stay, ventilation time and need for pharmacological inotropes, bleeding event. Surgical risk was calculated according to EuroSCORE II by evaluating following factors: age, gender, GFR_e, NYHA functional class, urgency and volume of the cardiac surgery, SPAP, LVEF, CCS functional class of angina, presence/absence of extracardiac arteriopathy, poor mobility, previous cardiac surgery, chronic obstructive pulmonary disease, active endocarditis, recent MI, DM on insulin treatment, critical state, thoracic aorta surgery (24, 25).

For post-op analysis surgery outcomes (alive vs died), complications (acute kidney injury (AKI), acute respiratory failure, chronic HF exacerbation) and arrhythmia (presence or absence) were used.

NT-proBNP assay

First sample was obtained in operation theatre prior to anesthesia. Second sampling was performed after 24 hours of operation in patients ventilated and intravenously supported by inotropes. Serum investigation was performed immediately after blood sampling in UPT – 3A analyzer, by electrochemiluminescence immunoassay (ECLIA) method, measurement units - pg/ml (Beijing Hotgen Biotech Co.LTD, China).

In adults a <300 pg/ml level of NT-proBNP could safely exclude likelihood of HF at any age. According to age aspects, for individuals aged 21-50yrs ≥ 450 pg/ml, 50-75yrs ≥ 900 pg/ml, elder 75yrs ≥ 1800 pg/ml levels confirm heart failure. The values between 300pg/ml and aforementioned levels remained as grey zone (26).

In pediatric patients to date, no clear standards and no normal ranges of NT-proBNP. Based on ROC curve some studies revealed the 598ng/l as a cut-off level of HF diagnosis in children (27).

We used as cut-off values 430pg/ml for adult cardiac surgery patients (19) and 250pg/ml for pediatric cardiac surgery patients (20).

Statistical analysis

Statistical analysis was performed using SPSS for Windows software (IBM, New York, USA).

Continuous variables are depicted in mean (standard deviation), median (min-max range), categorical variables as counts (percentage).

Chi-square test was used to assess differences for categorical variables, and Mann Whitney U test - for continuous variables. Nonparametric Spearman correlation analysis was used to assess correlation of NT-pro BNP level with baseline and clinical-perioperative variables. P value of less than 0.05 was accepted as significant value determination ($p < 0.05$).

Due to scarcity of data and small sample size we used only descriptive statistics for pediatric groups and Spearman correlation analysis of relationship of NTproBNP with clinical and perioperative variables.

Results

Adult cardiac surgery patients

Baseline population demographics

Patient population was categorized into two groups on the basis of preoperative NT-proBNP level. A 430pg/ml was accepted as the cutoff value for grouping in AG. Baseline clinical variables and their statistical results are highlighted in Table 1. Mean age 60.01 years. Age range: 34 to 72 years. All operations were performed in circumstance of CPB except the stent-graft implantation: CABG (n=15), aortic valve replacement (AVR) (n=4), mitral valve replacement (MVR) (n=3), aortic and mitral valve replacements (n=3), left atrial (LA) myxoma resection (n=1), stent-graft implantation (n=1).

According to age, sex, main diagnosis, surgery type, risk factors of HF and postoperative clinical outcomes no significant differences were found among groups. Significant results were obtained for BMI ranges and categorizations (slim ($<25\text{kg/m}^2$) and overweight or obese ($\geq 25\text{kg/m}^2$)): p-value was 0.03 and 0.017, respectively. More obese patients were in group of low NTproBNP level. There were significantly more patients with higher NYHA class in group of patients with high NTproBNP levels ($p<0.001$), especially more patients with class III ($p=0.03$).

Clinical and operative variables

All results are depicted in Table 2. Echocardiographic data regarding cardiac chamber values, LV systolic function and estimated SPAP did not show significant differences ($p>0.05$). Similar calculations were obtained from creatinine level, intraoperative and ICU parameters, namely, ventilation time (VT), CPB, ACCT, and duration of ICU stay. Nevertheless, significant differences were observed among groups by estimated glomerular filtration rate (GFR_e, $p=0.036$), frequency of renal function impairment (moderately impaired, $p=0.049$), need for inotropes in

early postop ($p=0.006$) and EuroSCORE II scale ($p=0.008$). Patients with high NTproBNP levels had markedly lower eGFR rate ($p=0.036$), higher number of patients with renal function impairment ($p=0.049$), and more patients in this group needed inotropic support (0.006). The patients with higher NTproBNP had higher surgical risk values compared to subjects with lower levels of NTproBNP ($p=0.008$). However, there were no differences in adverse outcomes and complications between groups. We also did not observe any bleeding events.

Correlations of NTproBNP with clinical and perioperative variables (Table 3)

NT proBNP levels positively, strongly and significantly correlated with EuroSCORE II, NYHA classes and GFR_e ($p=0.001$) with rho-values 0.73, 0.87 and -0.73, respectively. Significant results ($p<0.05$) were also found for the following parameters with corresponding rho-values: need for inotropes (0.62), renal function categories (0.72), EF obtained by Teichholz method (-0.47) and LV function categories (0.52).

NTproBNP was negatively correlated with BMI ranges (-0.73), meaning higher BNP levels in lower BMI values. But higher NTproBNP levels were related to lower values of EF (-0.47), and NTproBNP levels were positively correlated to systolic indices (categories $<40\%$ and $<50\%$ of EF) (rho 0.52, $p=0.01$).

Pediatric cardiac surgery patients

Baseline demographic indices

According to cut-off point of NT pro-BNP level PG was grouped into less or more than 250pg/ml. As can be seen from tables 4 and 5, only descriptive statistics of baseline clinical and operative parameters were highlighted. The mean age was 32.1 ± 30.3 months, median: 23 months, age range 14 days (0 months)-96 months (8 years).

Table 1. Baseline clinical characteristics of adult patients

Variable	Group 1 NT pro-BNP<430, pg/ml	Group 2 NT pro-BNP>430, pg/ml	p
Age, years			0.672
Mean (SD)	58.0 (12.17)	61.46 (6.32)	
Median (range)	60 (34-69)	62 (48-72)	
Sex, n (%)			0.604
Male	5 (71.4)	9 (60.0)	
Female	2 (28.6)	6 (40.0)	
Surgery, n (%)			0.928
CABG	4 (57.1)	8 (53.3)	
MVR	1 (14.3)	1 (6.7)	
AVR	1 (14.3)	3 (20)	
Myxoma resection	0	1 (6.7)	
AVR+MVR	1 (14.3)	2 (13.3)	
Diagnosis, n (%)			0.405
CAD	4 (57.1)	7 (46.7)	
RVHD	1 (14.3)	6 (40)	
BiAV	1 (14.3)	1 (6.7)	
Myxoma	0	1 (6.7)	
IE	1 (14.3)	0	
NYHA class, n (%)			<0.001
I	5 (71.4)	0	
II	2 (28.6)	5 (33.3)	
III	0	10 (66.7)	
BMI, kg/cm²			0.017
Mean (SD)	31.7 (3.15)	25.9 (3.8)	
Median (range)	30.7 (28.6-36.5)	25.5 (19.5-32.5)	
BMI categories, n (%)			0.06
<25	0	5 (50)	
25-30	2 (40)	4 (40)	
>30	3 (60)	1 (10)	
Dyslipidemia n (%)	3 (42.9)	9 (64.3)	0.35
Myocardial infarction, n (%)	1 (14.3)	5 (35.7)	0.306
Hypertension, n (%)	4 (57.1)	8 (66.7)	0.678
Diabetes mellitus, n (%)	6 (85.7)	11 (91.7)	0.683
Preoperative Arrhythmia, n (%)	0	3 (23.1)	0.23
Postoperative Arrhythmia, n (%)	0	8 (61.5)	0.03
Outcome, n (%)			0.746
Alive	6 (85.7)	12 (80)	
Died	1 (14.3)	3 (20)	
Complications, n (%)			0.469
AKI	0	1 (6.7)	
AKI+CHFe	0	1 (6.7)	
CHFe	1 (14.3)	0	
Resp.F+CHFe	0	1 (6.7)	
none	6 (85.7)	12 (80)	

AKI –acute kidney injury, AVR-aortic valve replacement, BMI-body-mass index, BiAV-bicuspid aortic valve, CABG-coronary artery by-pass grafting, CHFe-chronic heart failure exacerbation, MVR-mitral valve replacement, NTproBNP-N-terminal pro brain type natriuretic peptide, NYHA class-New-York Heart Association class, Resp.F-respiratory failure

Table 2. Clinical and operative variables in an adult cardiac surgery group

Variable	Group 1 NTproBNP<430 pg/ml	Group 2 NTproBNP>430 pg/ml	p
LVESD, mm			0.501
Mean (SD)	37.2 (10.6)	38.6 (7.6)	
Median (range)	33 (30-60)	39 (29-52)	
LVEDD, mm			0.626
Mean (SD)	54.4 (5.6)	53.9 (5.8)	
Median (range)	55 (43-60)	52 (46-63)	
EF Teichholz, %			0.156
Mean (SD)	63.4 (5.3)	56.6 (10.01)	
Median (range)	63 (57-70)	59 (34-69)	
LV function, n (%)			0.194
reduced EF<40%	0	1 (7.1)	
moderate reduced EF 40-50%	0	4 (28.6)	
preserved EF>50%	7 (100)	9 (64.3)	
EF Simpson, %			0.454
Mean (SD)	54.5 (4.13)	51.8 (6.3)	
Median (range)	54.5 (49-60)	54 (43-59)	
Creatinine, µmol/l			0.794
Mean (SD)	85.3 (19.4)	92.06 (21.9)	
Median (range)	85.3 (56.1-115.4)	86.3 (66.2-149.4)	
GFR_e, ml/min/1.73m²			0.036
Mean (SD)	111.4 (26.4)	77.9 (22.4)	
Median (range)	119.7 (78.8-137.7)	75.5 (55.4-132)	
RF, n (%)			0.049
Normal (>85 ml/min/1.73m ²)	4 (80)	3 (27.3)	
Moderate impairment (60-45 ml/min/1.73m ²)	1 (20)	8 (72.7)	
SPAP, mm Hg			0.187
Mean (SD)	34.6 (5.5)	39 (5.1)	
Median (range)	35 (28-40)	40 (30-50)	
PAH, n (%)			0.119
No (<30 mmHg)	2 (40)	1 (8.3)	
Moderate PAH (31-55 mmHg)	3 (60)	11 (91.7)	
ICU duration, days			0.817
Mean (SD)	3.8 (4.5)	2.6 (2.9)	
Median (range)	1.0 (1.0-1.13)	1.0 (1.0-1.10)	
CPB time, min			0.938
Mean (SD)	124.0 (93.7)	108.3 (58.6)	
Median (range)	84.5 (64.0-263.0)	103.0 (40.0-219.0)	
ACCT, min			0.643
Mean (SD)	88.2 (66.5)	74.3 (50.3)	
Median (range)	56.5 (52.0-188.0)	65.0 (29.0-180.0)	
Inotropes need, n (%)	2 (28.6)	13 (86.7)	0.006
Ventilation time, hours			0.84
Mean (SD)	38.5 (62.3)	9.7 (4.4)	
Median (range)	8.0 (6.0-132.0)	8.0 (4.0-19.0)	
EuroSCORE II			0.008
Mean (SD)	0.81 (0.19)	1.7 (0.5)	
Median (range)	0.75 (0.64-1.13)	1.83 (0.67-2.2)	

ACCT-aortic cross-clamp time, CPB-cardiac-pulmonary bypass, EF-left ventricular ejection fraction, GFR_e-estimated glomerular filtration rate, ICU-intensive care unit, LV – left ventricle, LVEDD-left ventricular end-diastolic diameter, LVESD-left ventricular end-systolic diameter, NTproBNP-N-terminal pro brain type natriuretic peptide, PAH-pulmonary arterial hypertension, RF-renal function, SPAP-systolic pulmonary artery pressure

Table 3. Correlation of NTproBNP with baseline, clinical and operative variables in adult group

Variable	Rho	p
EuroSCORE II	0.73	0.001
NYHA class	0.87	<0.001
Inotropes need	0.62	0.002
BMI	-0.73	0.002
EF Teichholz	-0.47	0.02
LV function	0.52	0.01
GFR _e	-0.73	0.001
Renal function	0.72	0.02

BMI – body mass index, GFR_e-estimated glomerular filtration rate, EF – ejection fraction, LV – left ventricle, NTproBNP-N-terminal pro brain type natriuretic peptide, NYHA class-New-York Heart Association class

Of 16 patients 9 are males (45%). The average NT pro-BNP level was 3602.5 (9094) pg/ml, min 217.9 pg/ml, max 35000pg/ml. Mean BSA: 0.48 (0.21m²), median: 0.45m² range: 0,21-0,91m². Patients were distributed according to underlying CHD as follows: ventricular septal defect (VSD, n=6), atrial septal defect (ASD, n=2), ASD+VSD (n=1), ASD+partial anomalous pulmonary venous return (PAPVR) (n=1), patent ductus arteriosus (PDA, n=1), Fallot tetralogy (TOF, n=2), complete atrioventricular septal defect (AVSDc, n=1), coarctation of aorta (CoA, n=1), pulmonary stenosis (PS, n=1). Performed surgical interventions with use of CPB are as follows: ASD closure (n=2), VSD correction (n=6), VSD+ASD closure (n=1), ASD closure and PAPVR repair (n=1), complete AVSD correction (n=1), radical correction of TOF (n=1), pulmonary valvulotomy (n=1). Surgical procedures without CPB were systemic-pulmonary shunting in TOF (n=1), PDA ligation (n=1), and aortic coarctation (CoA) correction (n=1). According to NYHA Ross classification children were distributed by: I (n=3), II (n=4) and III (n=9).

Clinical and operative variables

Echocardiographic indices: LVEDD: 29.07 (7.08) mm, LVESD: 17.28 (4.7) mm, LVEF: 72.1 (4.4)%. Systolic pulmonary artery pressure (SPAP): 56.8 (17.1)mmHg. Creatinine level: 38.2 (14.9) μmol/l. CPB time: 50.2 (21.4)min, ACCT: 26.0 (11.5) min, ICU duration: 2.07(1.6) days, ventilation time: 17.7 (34.8) h, Need for inotropic support: 12 (75%). Only single lethal case was observed during hospital stay due to exacerbation of CHF (0.62%).

Correlations

A non-parametric correlation represented statistically significant correlation of NTproBNP with following variables: age (p=0.038), BSA (p=0.002), left ventricular diastolic (p=0.024) and systolic diameters (p=0.053) (all are negatively correlated) (Table 6).

Table 4. Baseline clinical characteristics of pediatric patients

Variable	Group 1 NT pro-BNP<250, pg/ml	Group 2 NT pro-BNP>250, pg/ml	Number of cases
Age, months			14
Mean (SD)	52.0 (27.7)	26.7 (29.9)	
Median (range)	36 (36-84)	11 (1-96)	
Sex, n (%)			16
Male	1 (33.3)	6 (46.2)	
Female	2 (66.7)	7 (53.8)	
Surgery, n (%)			16
ASD closure		2 (15.3)	
VSD closure	3 (100)	3 (23.07)	
ASD+VSD correction		1 (7.6)	
PDA ligation		1 (7.6)	
AVSDc correction		1 (7.6)	
TOF radical correction		1 (7.6)	
MBTS		1 (7.6)	
CoA correction		1 (7.6)	
PV surgery		1 (7.6)	
Diagnosis, n (%)			16
ASD		2 (15.3)	
VSD	3 (100)	3 (23.07)	
ASD+VSD		1 (7.6)	
PDA		1 (7.6)	
ASD+PAPVR		1 (7.6)	
AVSDc		1 (7.6)	
PS		1 (7.6)	
TOF		2 (15.3)	
CoA		1 (7.6)	
NYHA Ross, n (%)			16
I	1 (33.3)	2 (15.3)	
II	1 (33.3)	3 (23.07)	
III	1 (33.3)	8 (61.5)	
BSA, m2			14
Mean (SD)	0.7 (0.17)	0.42 (0.18)	
Median (range)	0.65 (0.59-0.91)	0.37 (0.21-0.80)	
Outcome, n (%)			16
Alive	3 (100)	12 (92.3)	
Died	0	1 (7.6)	
Complications, n (%)			16
CHFe	0	1 (7.6)	
None	3 (100)	12 (92.3)	

ASD—atrial septal defect, AVSDc-complete atrio-ventricular septal defect, BSA-body surface area, CHFe-exacerbation of chronic heart failure, CoA- coarctation of aorta, MBTS-modified Blalock-Tausig shunt, NTproBNP-N-terminal pro brain type natriuretic peptide, NYHA Ross-Modified Ross New-York Heart Association functional class, PAPVR-partial anomalous pulmonary venous return, PDA-patent ductus arteriosus, PS-pulmonary stenosis, PV – pulmonary valvotomy, TOF-Tetralogy of Fallot, VSD-ventricular septal defect

Table 5. Clinical and operative variables in pediatric patients

Variable	Group 1 NTproBNP<250 pg/ml	Group 2 NTproBNP>250 pg/ml	Number of cases
LVESD, mm			14
Mean (SD)	20.0 (2.6)	16.5 (4.9)	
Median (range)	19 (18-23)	16 (10-25)	
LVEDD, mm			14
Mean (SD)	33.6 (5.5)	27.8 (7.1)	
Median (range)	34 (28-39)	28 (16-41)	
EF Teichholz, %			14
Mean (SD)	71.0 (5.5)	72.4 (4.3)	
Median (range)	72 (65-76)	73 (64-78)	
Creatinine, µmol/l			14
Mean (SD)	47.5 (14.4)	35.7 (14.7)	
Median (range)	49.4 (32.2-60.9)	37.1 (17.6-58.6)	
SPAP, mm Hg			10
Mean (SD)	47.5 (3.5)	59.1 (18.6)	
Median (range)	47.5 (45-50)	57.5 (30-85)	
ICU duration, days			14
Mean (SD)	1.3 (0.57)	2.2 (1.7)	
Median (range)	1 (1.0-2.0)	1.0 (1.0-1.6)	
CPB time, min			11
Mean (SD)	64.3 (28.5)	45.0 (17.5)	
Median (range)	75.0 (32.0-86.0)	45.0 (19.0-73.0)	
ACCT, min			11
Mean (SD)	32.6 (18.4)	23.5 (8.1)	
Median (range)	28.0 (17.0-53.0)	24.0 (11.0-34.0)	
Inotropes need, n (%)	1 (33.3)	11 (84.6)	16
Ventilation time, hours			11
Mean (SD)	7.6 (1.5)	20.5 (39.2)	
Median (range)	8.0 (6.0-9.0)	8.0 (2.0-138.0)	

ACCT-aortic cross-clamp time, CPB-cardiac-pulmonary bypass, EF-left ventricular ejection fraction, ICU-intensive care unit, LVEDD-left ventricular end-diastolic diameter, LVESD-left ventricular end-systolic diameter, NTproBNP-N-terminal pro brain type natriuretic peptide, SPAP-systolic pulmonary artery pressure

Table 6. Correlation of NTpro-BNP with clinical and operative variables in pediatric patients

Variable	rho	p
Age	-0.55	0.038
BSA	-0.75	0.002
LV EDD	-0.59	0.024
LV ESD	-0.52	0.053

BSA-body surface area, LVEDD-left ventricular end-diastolic diameter, LVESD-left ventricular end-systolic diameter, NTproBNP-N-terminal pro brain type natriuretic peptide

Discussion

Preliminary results of our study in adult patients who underwent cardiac surgery demonstrated that preoperative NT pro-BNP levels are elevated in patients with advanced NYHA classes, LV dysfunction, higher surgical risk, worsened kidney function and need of inotropic support. In addition to, significant positive correlation was found between NT pro-BNP level and EuroSCORE II, a prognostic scoring index which defines the complexity of surgery and patients' cardiac surgical risk; renal function categories, whether normal or moderately impaired; CHF NYHA functional class grading, from mild symptoms to limited daily activities; systolic function categories on the basis of LVEF; as well as need for inotropic support in early post-op. A negative correlation was observed by BMI, EF and GFR levels. However, we did not observe difference in surgery outcomes due presumably to small sample size.

Previous studies revealed the good correlation of LV global systolic dysfunction and symptom severity with NT-pro-BNP by means of LVEF with cut-off level of 50% (28). Nevertheless, recent findings advocate the relationships of BNP molecules with novel indices of global systolic function. According to Ablasser et al, LV global longitudinal strain was significantly correlated with NT-proBNP level in patients with ischemic (Pearson $r=0.266$, $p=0.049$) and non-ischemic etiology of heart failure (Pearson $r=0.434$, $p=0.034$) but not the ejection fraction (29). Next studies necessitate the evaluation of left ventricular systolic function with speckle-tracking derived methods which are considered as more reliable and promising options.

Several studies including large trials represented negative correlations between natriuretic peptides and BMI. According to these studies low NT pro-BNP levels were detected both in overweight (BMI=25-30kg/m²) and obese (BMI≥30kg/m²) subjects. Furthermore, these investigations defined similar levels of NT pro-BNP levels as the cutoffs, in PRIDE-study which included 204 subjects was chosen 900pg/ml, in another trial with 1103 study population considered 986ng/l (30, 31). Recent evidence supports the involvement of posttranslational processing of pro-BNP molecules, specifically the augmented glycosylation of threonine in 71st position of pro-BNP and consequent decreasing in concentration of NT pro-BNP molecules in overweight and obese patients (32). Despite the small sample size (n=28) our patients also represented negative correlation with statistically significant results ($\rho=-0.73$, $p=0.002$) as in aforementioned literature reviews.

A univariate analysis showed that renal dysfunction was closely correlated with NT pro-BNP level in assessment of the 30-day mortality (18). According to Su et al, NT-pro-BNP was found as an independent risk factor for development of acute kidney injury requiring renal replacement therapy in early post-op of cardiac surgery patients (33). Both continuous (GFR_e) and categorical (renal impairment) variables in our study showed significant results among adult subgroups, as well as significant negative correlation of eGFR and positive correlation with renal impairment severity with NTproBNP level, thus confirming previous studies.

Controversial findings exist in literature regarding to correlation of NT-pro-BNP and inotropic support in early post-op. According to Oztekin et al., no relationship was found between NT-proBNP level and need for inotropic support in patients underwent isolated CABG-surgery (34). In pediatric cardiac surgery, perioperative NT-pro-BNP accurately determines the prognosis of post-op outcome including duration of mechanical ventilation, ICU stay and inotropic therapy (35).

Recent findings revealed the reliability and simplicity of pre-op NT-pro-BNP in predicting cardiac surgery risk or assisting the EuroSCORE II grading system in a large body of evidence (36, 37). Our results are in agreement with those studies showing strong positive correlation with EuroSCORE II.

Study limitations

The main limitations of this preliminary study are small sample size and retrospective design. Results will be validated and reported in further larger sample size study.

Conclusion

Preoperative NT-pro-BNP level in our cohort of adult patients is related to higher NYHA class, LV dysfunction, renal impairment and consequently higher requirement for inotropic support after surgery and is related with prognostic indicator as EuroSCORE II independently of type of surgery. There is a negative relation with BMI. Further study is needed to support our findings in adult and pediatric patients.

Peer-review: External and internal

Conflict of interest: None to declare

Authorship: I.A.A., T.Z.K., D.A.A., A.A.Zh., D.E.Z., K.T.T., G.N.N. are equally contributed to the study and preparation of article

Acknowledgments and funding: None to declare

References

1. Mareev VY, Fomin IV, Ageev FT, Begrambekova YL, Vasyuk YA, Garganeeva AA, Gendlin GE, et al. Russian Heart Failure Society, Russian Society of Cardiology. Russian Scientific Medical Society of Internal Medicine Guidelines for Heart failure: chronic (CHF) and acute decompensated (ADHF). Diagnosis, prevention and treatment. *Kardiologiia* 2018; 58: 8-158.
2. Mareev VY, Ageev FT, Arutyunov GP, Korolev AV, Revishvili AS. Diagnosis and treatment of chronic heart failure. Working group on guideline preparation 2010.
3. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS, et al. 2016 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure. The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J* 2016; 37: 2129-200.
4. Hunt SA, Abraham WT, Chin MH, Feldman AM, Francis GS, Ganiats TG, et al. American College of Cardiology; American Heart Association Task Force on Practice Guidelines; American College of Chest Physicians; International Society for Heart and Lung Transplantation; Heart Rhythm Society. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure): developed in collaboration with the American College of Chest Physicians and the International Society for Heart and Lung Transplantation: endorsed by the Heart Rhythm Society. *Circulation* 2005; 112: e154-235. DOI: 10.1161/CIRCULATIONAHA.105.167586
5. Lindenfeld J, Albert NM, Boehmer JP, Collins SP, Ezekowitz JA, Givertz MM, et al. HFSA 2010 comprehensive heart failure practice guideline. *J Card Fail* 2010; 16: e1-194. doi: 10.1016/j.cardfail.2010.04.004.
6. Bayés-Genís A. The circulating NTproBNP level, a new biomarker for the diagnosis of heart failure in patients with acute shortness of breath). *Rev Esp Cardiol* 2005; 58: 1142-4.
7. Yaffee DW, Williams MR. Cardiovascular surgery in the elderly. *Semin Thorac Cardiovasc Surg* 2016; 28: 741-7. doi:10.1053/j.semtcvs.2016.08.007
8. Parissis H. Cardiac surgery: what the future holds? *J Cardiothorac Surg* 2011; 6: 93. doi:10.1186/1749-8090-6-93.
9. Qu J, Liang H, Zhou N, Lijuan L, Wang Y, Jianbin L, et al. Perioperative NT-proBNP level: Potential prognostic markers in children undergoing congenital heart disease surgery. *J Thorac Cardiovasc Surg* 2017; 154: 631-40. doi:10.1016/j.jtcvs.2016.12.056.
10. Hutfless R, Kazanegra R, Madani M, Bhalla MA, Tata AT, Chen A, et al. Utility of B-type natriuretic peptide in predicting postoperative complications and outcomes in patients undergoing heart surgery. *Journal of the American College of Cardiology* 2004; 43: 1873-9. <https://doi.org/10.1016/j.jacc.2003.12.048>
11. Soo-Hoo S, Nemeth S, Baser O, Argenziano M1, Kurlansky P; East meets West: the influence of racial, ethnic and cultural risk factors on cardiac surgical risk model performance. *Heart Asia* 2018; 10: e010995. doi: 10.1136/heartasia-2017-010995.
12. Yasue H, Yoshimura M, Sumida H, Kikuta K, Kugiyama K, Jougasaki M, et al. Localization and mechanism of secretion of B-type natriuretic peptide in comparison with those of A-type natriuretic peptide in normal subjects and patients with heart failure. *Circulation* 1994; 90: 195–203.
13. Cowie MR, Struthers AD, Wood DA, Coats AJ, Thompson SG, Poole-Wilson PA, et al. Value of natriuretic peptides in assessment of patients with possible new heart failure in primary care. *Lancet* 1997; 350: 1349-53.
14. McDonagh TA, Robb SD, Murdoch DR, Morton JJ, Ford I, Morrison CE, et al. Biochemical detection of left-ventricular systolic dysfunction. *Lancet* 1998; 351: 9-13.
15. Johannes Steiner, Maya Guglin; BNP or NTproBNP? A clinician's perspective. *International Journal of Cardiology* 2008; 129: 5-14.
16. Duceppe E, Patel A, Chan MT, Berwanger O, Ackland G, Kavsak PA, et al. Preoperative N-terminal pro-B-type natriuretic peptide and cardiovascular events after noncardiac surgery: a cohort study. *Ann Intern Med* 2020; 172: 96–104. doi: 10.7326/M19-2501.
17. Jogia, PM, Kalkoff M, Sleigh JW, Bertinelli A, La Pine M, Richards AM, et al. NT-Pro BNP Secretion and Clinical Endpoints in Cardiac Surgery Intensive Care Patients. *Anaesthesia and Intensive Care* 2007; 35: 363-9. Doi: 10.1177/0310057X0703500307.

18. Liu H, Wang C, Liu L, Zhuang Y, Yang X, Zhang Y. Perioperative application of N-terminal pro-brain natriuretic peptide in patients undergoing cardiac surgery. *J Cardiothorac Surg* 2013; 8: 1. doi:10.1186/1749-8090-8-1.
19. Schachner T, Wiedemann D, Fetz H, Laufer G, Kocher A, Bonaros N. Influence of preoperative serum N-terminal pro-brain type natriuretic peptide on the postoperative outcome and survival rates of coronary artery bypass patients. *Clinics* 2010; 65: 1239-45.
20. Zheng H, Li L, Liang H, Qu J, Li J, Cui Y, Chen X. Normalization of N-terminal pro-B-type natriuretic peptide after cardiac surgery among children with tetralogy of Fallot. *Am J Cardiol* 2018; 122: 2125-30. doi: 10.1016/j.amjcard.2018.08.052.
21. Specifications Manual for Joint Commission National Quality Measures (v2018A) Discharges 07-01-18 (3Q18) through 12-31-18 (4Q18). Available from: <https://manual.jointcommission.org/releases/TJC2018A/DataElem0439.html>
22. Ross RD, Bollinger RO, Pinsky WW. Grading the severity of congestive heart failure in infants. *Pediatr Cardiol* 1992; 13: 72-5.
23. KDIGO 2012 clinical practice guideline for evaluation and management of CKD. *Kidney Int Suppl* 2013; 3: 1-150. doi:10.1038/kisup.2012.72.
24. Nashef SA, Roques F, Sharples LD, Nilsson J, Smith C, Goldstone AR, et al. EuroSCORE II. *Eur J Cardiothorac Surg* 2012; 41: 734-45.
25. Euroscore II. Available from: URL: <https://www.mdcalc.com/european-system-cardiac-operative-risk-evaluation-euroscore-ii>
26. Januzzi JL, Camargo CA, Anwaruddin S, Baggish AL, Chen AA, Krauser DG, et al. The N-terminal Pro-BNP investigation of dyspnea in the emergency department (PRIDE) study. *Am J Cardiol* 2005; 95: 948-54.
27. Lin C, Zeng X, Jiang S, Wu T, Wang J, Zhang J, et al. Role of the NT-proBNP level in the diagnosis of pediatric heart failure and investigation of novel combined diagnostic criteria. *Exp Ther Med* 2013; 6: 995-9. <https://doi.org/10.3892/etm.2013.1250>.
28. O'Donoghue M, Chen A, Baggish AL, Anwaruddin S, Krauser DG, Tung R, et al. The effects of ejection fraction on N-terminal proBNP and BNP levels in patients with acute CHF: Analysis from the ProBNP Investigation of Dyspnea in the Emergency Department (PRIDE) Study. *J Card Fail* 2005; 11: S9-S14. doi: 10.1016/j.cardfail.2005.04.011.
29. Ablasser K, Lewinski DV, Kolesnik E, Gangl M, Kattinig L, et al. Comparison of global longitudinal strain and ejection fraction in correlation to NT-proBNP in ischemic and non-ischemic heart failure. *Eur Heart J* 2019; 40: ehz748.0938. doi: 10.1093/eurheartj/ehz748.0938
30. Wirth J, Buijsse B, di Giuseppe R, Fritsche A, Hense HW, Westphal S, et al. Relationship between N-terminal pro-brain natriuretic peptide, obesity and the risk of heart failure in middle-aged German adults. *PLoS One* 2014; 9: e113710. doi: 10.1371/journal.pone.0113710.
31. Krauser DG, Lloyd-Jones DM, Chae CU, Cameron R, Anwaruddin S, Baggish AL, et al. Effect of body mass index on natriuretic peptide levels in patients with acute congestive heart failure: a ProBNP Investigation of Dyspnea in the Emergency Department (PRIDE) substudy. *Am Heart J* 2005; 149: 744-50. doi: 10.1016/j.ahj.2004.07.010.
32. Lewis LK, Raudsepp SD, Prickett TCR, Yandle TG, Pemberton CJ, Richards AM. ProBNP processing is decreased by obesity in patients with heart failure. *ATM* 2020; 8: doi: 10.21037/atm.2019.12.81.
33. Su Y, Hou JY, Zhang YJ, Ma GG, Hao GW, Luo JC, et al. Serum N-terminal pro-B-type natriuretic peptide predicts mortality in cardiac surgery patients receiving renal replacement therapy. *Front Med (Lausanne)*. 2020; 7: 153. doi: 10.3389/fmed.2020.00153.
34. Öztekin A, Memetoglu M, Kutlu R, Tekin A, Erbasan O, Arslan U, et al. The predictive value of preoperative serum NTproBNP levels for the need for inotropic support in the postoperative period in patients undergoing coronary artery bypass grafting. *Cardiovasc J* 2017; 9: 90-6. doi: 10.3329/cardio.v9i2.32419.
35. Qu J, Liang H, Zhou N, Li L, Wang Y, Li J, et al. Perioperative NT-proBNP level: Potential prognostic markers in children undergoing congenital heart disease surgery. *J Thorac Cardiovasc Surg* 2017; 154: 631-40.
36. Brynildsen J, Petaja L, Pettila V, Nygard S, Vaara ST, Linko R et al. The predictive value of NT-proBNP and hs-TnT for risk of death in cardiac surgical patients. *Clin Biochem* 2018; 53: 65-71.
37. Holm J, Vidlund M, Vanky F, Friberg O, Hakanson E, Walther S, et al. EuroSCORE II and N-terminal pro-B-type natriuretic peptide for risk evaluation: an observational longitudinal study in patients undergoing coronary artery bypass graft surgery. *BJA: Br J Anaesth* 2014; 113: 75-82. doi: 10.1093/bja/aeu