

A rare case of bacterial infective endocarditis caused by *Streptococcus alactolyticus*

Nedim Cekmen, *Oben Baysan, Emine Disbudak, Ceren Gunt

Intensive Care Unit, *Cardiology Department, Guven Hospital, Ankara, Turkey

Abstract

Background: *Streptococcus alactolyticus* is a rarely isolated bacterium, which classified under DNA cluster IV of the *S. Bovis/S. equinus* complex. Infections, especially infective endocarditis, caused by *Strep. alactolyticus* are very rare in humans.

Case Report: We describe a case of *Streptococcus alactolyticus* bacteremia complicated by infective endocarditis. A 64-year-old male with a previous history of coronary artery bypass grafting applied to our cardiology outpatient clinic with complaints of dyspnea, fever, confusion and an apical holosystolic murmur. He was admitted to the intensive care unit. Transthoracic and transesophageal echocardiography showed the presence vegetation on the aortic valve. *S. alactolyticus* was detected on serial blood cultures. The patient was first treated with intensive antimicrobial therapy, and then underwent mitral and aortic valve replacements with uneventful follow-up.

Conclusion: *Streptococcus alactolyticus* infective endocarditis has only been reported previously in one patient. More information is certainly needed for diagnosis and treatment of patients infected with *Streptococcus alactolyticus*.

Key words: *Streptococcus alactolyticus*, endocarditis

(Heart, Vessels and Transplantation 2019; 3: 109-113. doi: 10.24969/hvt.2019.133)

Introduction

Streptococcus alactolyticus is a particular subspecies grouped under *streptococcal bovis/Streptococcal equinus complex* (SBSEC) (1, 2). This complex contains *non-beta hemolytic streptococcus* and *Lancefield group D streptococci*. They are commensal colonizers of human and animal gastrointestinal tract and act as opportunistic pathogens (3).

S. alactolyticus has been isolated from intestinal flora of pigs, chicken, cows, pigeons and dogs (4, 5). Human infections caused by *S. alactolyticus* are extremely rare. Herein, we describe a rare case of bacterial infective endocarditis (IE) caused by *S. alactolyticus*.

Case report

A 64-year-old male was transferred to our emergency service by his family with difficulty of breathing, tachycardia, fever

and confusion. He had previous history of coronary artery bypass graft surgery one month before. On admission, his vital signs were as follows high respiratory rate (32 breaths/min), temperature 38.6 C°, blood pressure 150/90 mmHg, heart rate 132 beats/min, and SpO2 67 %. He was urgently transferred to the intensive care unit (ICU). He had diminished respiratory sounds with crepitating rales bilaterally. We detected a grade 3/6 an apical holosystolic murmur on his left precordium. Glasgow coma score was 9, Acute Physiology and Chronic Health Evaluation score (APACHE II) was 18 and his Multiple Organ Dysfunction Score (MODS) was 6. Arterial blood gas analysis showed metabolic acidosis and hypoxemia: (bicarbonate 17.2 mmol/L, pH 7.14, PaCO2 35 mm Hg, and PaO2 47 mm Hg while on nasal oxygen at 6 L/min).

Address for Correspondence: Oben Baysan, Guven Hospital, Simsek Street No:29 Kavaklidere, Ankara, Turkey Mobile: +905322718225
Email: obbaysan@gmail.com

Received: 30.05.2019 **Accepted:** 7.06.2019

Copyright©2019 Heart, Vessels and Transplantation

Blood cultures were taken. He was intubated and mechanically ventilated. Laboratory analyses showed high leukocyte count (23,400/L) with neutrophil dominance (83%), erythrocyte sedimentation rate 62 mm/h, C-reactive protein 135 mg/L (reference range: 0-5 mg/L), procalcitonin 2.4 ng/L, and creatinine 1.4 mg/dL. Immediate bedside transthoracic echocardiography revealed a suspicious mass on the aortic valve with severe aortic regurgitation. There was also a moderate degree mitral regurgitation. Same day urgent transesophageal echocardiogram (TEE) confirmed presence of a large mobile vegetation (23x15 mm) on aortic valve associated with severe aortic regurgitation (Fig. 1 and 2) and moderate degree mitral regurgitation (Fig. 3). Empiric antimicrobial therapy with vancomycin, gentamicin, and penicillin G was initiated. His general condition showed a noticeable improvement within 48 hours. The *group D streptococci* were identified from blood cultures as *S. alactolyticus* by Vitek 2 (bioMerieux Vitek, Inc, Hazelwood, St. Louis, MO, USA). It was susceptible to penicillin. The second set of blood cultures were also positive for

S. alactolyticus. Vancomycin was discontinued and intravenous penicillin G (3,000,000 U every 4 hr) and gentamicin (63 mg every 8 hr) were continued. With that therapy, he showed persistent clinical improvement. His subsequent blood cultures were negative. The patient was extubated on the 10th day of his intubation. The neurologist's consultation including brain magnetic resonance imaging (MRI) was unremarkable for the presence of cerebral embolism. We recommended to the patient early surgical intervention. He agreed with the proposal and underwent valvular heart surgery with bioprosthetic aortic and mitral valve replacements. Based on the *S. alactolyticus* sensitivities, antimicrobial therapy was deescalated to ceftriaxone, which was planned to continue for another six-weeks. The patient recovered completely and was discharged on the 44th day of hospitalization. He was well on follow-up 8 weeks after discharge. He was also informed about an outpatient gastroenterology consultation for possible colonoscopy procedure.




Figure 1. A large aortic valve vegetation on TEE image (Mid esophageal 134 degree)
TEE- transesophageal echocardiography




Figure 2. Severe aortic regurgitation caused by infective endocarditis.




Figure 3. Moderate mitral regurgitation

Discussion

S. Bovis/S. equinus complex bacteria contains both good strains such as *S. gallolyticus subsp. macedonicus* and *S. lutetiensis* ingested in daily food products and pathogenic strains (6). They inhabit gastrointestinal tract of animals and humans (7). Colon and ileum are major colonization sites for *S. Bovis/S. equinus complex* and they have a well-known association with colorectal cancer (8, 9). Bacteremia and infective endocarditis are frequently encountered diseases for human SBSEC infection.

Geographical location and development status of a country determine distribution of main pathogens causing IE. Indeed, *Staphylococcus aureus* is still the most frequently detected bacteria for all IE cases in developed countries (31%) followed by *viridans group streptococci* (17%), *coagulase negative staphylococci* (11%), *enterococci* (10%) and finally *S. Bovis/S. equinus complex (SBSEC)* (6%) (10). In contrast, *Streptococcus* still predominates in low to middle income countries (11, 12). Interestingly, the percentage of SBSEC-associated IE within all cases of streptococcal IE increased from 10.9% to 23.3% from 1995 to 2005 (6). Within SBSEC bacteria *S. gallolyticus subsp. gallolyticus* is a most prominent agent in IE cases across Europe. Another bacteria in this complex, *S. gallolyticus subsp. Pasteurianus*, is also the leading agent in Asia (13).

Human reports of *S. alactolyticus* infection are exceedingly rare. *S. alactolyticus* was reported as a causative agent in a patient with IE complicated by septic emboli (14). Another fatal case of fulminant neonatal sepsis caused by this pathogen was also reported (15).

Our patient presented with *S. alactolyticus bacteriemia* and aortic valve endocarditis. Aortic valve is the least frequently involved heart valve in SBSEC IE (16). In contrast, there is a tendency for SBSEC bacteria to affect multiple heart valves including mitral and prosthetic valves (16). The reported frequency of embolic events in SBSEC IE ranged from 9% to 55% (17, 18). Fortunately, the patient in our case had no embolic event confirmed by a neurologic exam and brain MRI. That could be related to relatively early diagnosis and prompt starting of antimicrobial therapy.

Conclusion

There is unmet need for better clarification of infective endocarditis caused by SBSEC bacteria. An umbrella term SBSEC contains many subspecies. The new SBSEC taxonomical grouping (1) brings new challenges and opportunities to clinicians. More reliable estimates of SBSEC related IE frequency and incidence could be obtained via the correct identification of SBSEC bacteria in each IE case, which provides better clarification of SBSEC IE epidemiology.

Peer-review: internal and external

Conflict of interest: None to declare

Authorship: N.C., O.B., E.D., C.G. equally contribute to case management and preparation of case report

Acknowledgement and funding: None to declare

References

- Schlegel L. Reappraisal of the taxonomy of the *Streptococcus bovis*/*Streptococcus equinus* complex and related species: description of *Streptococcus gallolyticus subsp. gallolyticus subsp. nov.*, *S. gallolyticus subsp. macedonicus subsp. nov.* and *S. gallolyticus subsp. pasteurianus subsp. nov.* Int J System Evolut Microbiol 2003; 53 :631.
- Facklam R. What happened to the streptococci: overview of taxonomic and nomenclature changes. Clin Microbiol Rev 2002;15: 613.
- Dekker JP, Lau AF, Kraft CS. An update on the *Streptococcus bovis* group: classification, identification, and disease associations. J Clin Microbiol 2016; 54: 1694.
- Farrow JAE, Kruse J, Phillips BA, Bramley A, Collins MD. Taxonomic studies on *Streptococcus bovis* and *Streptococcus equinus*: Description of *Streptococcus alactolyticus sp. nov.* and *Streptococcus saccharolyticus sp. nov.* System Appl Microbiol 1984;5:467-82.
- Rinkinen ML, Koort JMK, Ouwehand AC, Westermarck E, Björkroth KJ. *Streptococcus alactolyticus* is the dominating culturable lactic acid bacterium species in canine jejunum and feces of four fistulated dogs. FEMS Microbiol Letters 2004; 230: 35-9.
- Jans C, Meile L, Lacroix C, Stevens MJA. Genomics, evolution, and molecular epidemiology of the *Streptococcus bovis*/*Streptococcus equinus* complex (SBSEC). Infection, genetics and evolution. J Mol Epidemiol Evolut Genet Infect Dis 2015; 33: 419-36.
- Jans C, Boleij A. The road to infection: host-microbe interactions defining the pathogenicity of *Streptococcus bovis*/*Streptococcus equinus* complex members. Front Microbiol 2018; 9: 603.
- Dahmus JD, Kotler DL, Kastenber DM, Kistler CA. The Gut microbiome and colorectal cancer: a review of bacterial pathogenesis. J Gastrointest Oncol 2018; 9: 769-77.
- Boleij A, Roelofs R, Schaeps RMJ, Schülin T, Glaser P, Swinkels DW, et al. Increased exposure to bacterial antigen RpL7/L12 in early stage colorectal cancer patients. Cancer 2010; 116: 4014-22.
- Murdoch DR, Corey GR, Hoen B, Miró JM, Fowler VG, Bayer AS, et al. Clinical presentation, etiology, and outcome of infective endocarditis in the 21st century: the International Collaboration on Endocarditis-Prospective Cohort Study. Arch Intern Med 2009; 169: 463-73.
- Mirabel M, Rattanavong S, Frichitthavong K, Chu V, Kesone P, Thongsith P, et al. Infective endocarditis in the Lao PDR: clinical characteristics and outcomes in a developing country. Int J Cardiol 2015;180:270-3.
- Math RS, Sharma G, Kothari SS, Kalaivani M, Saxena A, Kumar AS, et al. Prospective study of infective endocarditis from a developing country. Am Heart J 2011; 162: 633-8.

13. Lee RA, Woo PCY, To APC, Lau SKP, Wong SSY. Geographical difference of disease association in *Streptococcus bovis* bacteraemia. *J Med Microbiol* 2003; 52: 903-8.
14. Almeida P, Railsback J, Gleason JB. A rare case of *Streptococcus alactolyticus* infective endocarditis complicated by septic emboli and mycotic left middle cerebral artery aneurysm. *Case Rep Infect Dis* 2016; 2016: 9081352.
15. Toepfner N, Shetty S, Kunze M, Orłowska-Volk M, Krüger M, Berner M, et al. Fulminant neonatal sepsis due to *Streptococcus alactolyticus* - A case report and review. *APMIS* 2014; 122: 654-6.
16. Hoen B, Chirouze C, Cabell CH, Selton-Suty C, Duchêne F, Olaison L, et al. Emergence of endocarditis due to group D streptococci: findings derived from the merged database of the International Collaboration on Endocarditis. *Eur J Clin Microbiol Infect Dis* 2005; 24: 12-6.
17. Kupferwasser I, Darius H, Müller AM, Mohr-Kahaly S, Westermeier T, Oelert H, et al. Clinical and morphological characteristics in *Streptococcus bovis* endocarditis: a comparison with other causative microorganisms in 177 cases. *Heart* 1998; 80: 276-80.
18. Pergola V, Di Salvo G, Habib G, Avierinos JF, Philip E, Vailloud JM, et al. Comparison of clinical and echocardiographic characteristics of *Streptococcus bovis* endocarditis with that caused by other pathogens. *Am J Cardiol* 2001; 88: 871-5.